

Improved Lower Bounds on the Compatibility of Multi-State Characters^{*}

Brad Shalters, Sudheer Vakati, and David Fernández-Baca

Department of Computer Science, Iowa State University, Ames, IA 50011, USA
{shalters,svakati,fernande}@iastate.edu

Abstract. We study a long standing conjecture on the necessary and sufficient conditions for the compatibility of multi-state characters: There exists a function $f(r)$ such that, for any set C of r -state characters, C is compatible if and only if every subset of $f(r)$ characters of C is compatible. We show that for every $r \geq 2$, there exists an incompatible set C of $\lfloor \frac{r}{2} \rfloor \cdot \lceil \frac{r}{2} \rceil + 1$ r -state characters such that every proper subset of C is compatible. Thus, $f(r) \geq \lfloor \frac{r}{2} \rfloor \cdot \lceil \frac{r}{2} \rceil + 1$ for every $r \geq 2$. This improves the previous lower bound of $f(r) \geq r$ given by Meacham (1983), and generalizes the construction showing that $f(4) \geq 5$ given by Habib and To (2011). We prove our result via a result on quartet compatibility that may be of independent interest: For every integer $n \geq 4$, there exists an incompatible set Q of $\lfloor \frac{n-2}{2} \rfloor \cdot \lceil \frac{n-2}{2} \rceil + 1$ quartets over n labels such that every proper subset of Q is compatible. We contrast this with a result on the compatibility of triplets: For every $n \geq 3$, if R is an incompatible set of more than $n - 1$ triplets over n labels, then some proper subset of R is incompatible. We show this upper bound is tight by exhibiting, for every $n \geq 3$, a set of $n - 1$ triplets over n taxa such that R is incompatible, but every proper subset of R is compatible.

Keywords: phylogenetics, quartet compatibility, character compatibility, perfect phylogeny

1 Introduction

The multi-state character compatibility (or perfect phylogeny) problem is a basic question in computational phylogenetics [18]. Given a set C of characters, we are asked whether there exists a phylogenetic tree that displays every character in C ; if so, C is said to be compatible, and incompatible otherwise. The problem is known to be NP-complete [3,21], but certain special cases are known to be polynomially-solvable [1,6,12,14,15,16,19]. See [8] for more on the perfect phylogeny problem.

In this paper we study a long standing conjecture on the necessary and sufficient conditions for the compatibility of multi-state characters.

^{*} This work was supported in part by the National Science Foundation under grants CCF-1017189 and DEB-0829674.

Conjecture 1 *There exists a function $f(r)$ such that, for any set C of r -state characters, C is compatible if and only if every subset of $f(r)$ characters of C is compatible.*

If Conjecture 1 is true, it would follow that we can determine if any set C of r -state characters is compatible by testing the compatibility of each subset of $f(r)$ characters of C , and, in case of incompatibility, output a subset of at most $f(r)$ characters of C that is incompatible.

A classic result on binary character compatibility shows that $f(2) = 2$; see [5,7,12,17,18]. In 1975, Fitch [9,10] gave an example of a set C of three 3-state characters such that C is incompatible, but every pair of characters in C is compatible, showing that $f(3) \geq 3$. In 1983, Meacham [17] generalized this example to r -state characters for every $r \geq 3$, demonstrating a lower bound of $f(r) \geq r$ for all r ; see also [16]. A recent breakthrough by Lam, Gusfield, and Sridhar [16] showed that $f(3) = 3$. While the previous results could lead one to conjecture that $f(r) = r$ for all r , Habib and To [13] recently disproved this possibility by exhibiting a set C of five 4-state characters such that C is incompatible, but every proper subset of the characters in C are compatible, showing that $f(4) \geq 5$. They conjectured that $f(r) \geq r + 1$ for every $r \geq 4$.

The main result of this paper is to prove the conjecture stated in [13] by giving a quadratic lower bound on $f(r)$. Formally, we show that for every integer $r \geq 2$, there exists a set C of r -state characters such that all of the following conditions hold.

1. C is incompatible.
2. Every proper subset of C is compatible.
3. $|C| = \lfloor \frac{r}{2} \rfloor \cdot \lceil \frac{r}{2} \rceil + 1$.

Therefore, $f(r) \geq \lfloor \frac{r}{2} \rfloor \cdot \lceil \frac{r}{2} \rceil + 1$ for every $r \geq 2$.

Our proof relies on a new result on quartet compatibility which we believe is of independent interest. We show that for every integer $n \geq 4$, there exists a set Q of quartets over a set of n labels such that all of the following conditions hold.

1. Q is incompatible.
2. Every proper subset of Q is compatible.
3. $|Q| = \lfloor \frac{n-2}{2} \rfloor \cdot \lceil \frac{n-2}{2} \rceil + 1$.

This represents an improvement over the previous lower bound on the maximum cardinality of such an incompatible set of quartets of $n - 2$ given in [21].

We contrast our result on quartet compatibility with a result on the compatibility of triplets: For every $n \geq 3$, if R is an incompatible set of triplets over n labels, and $|R| > n - 1$, then some proper subset of R is incompatible. We show this upper bound is tight by exhibiting, for every $n \geq 3$, a set of $n - 1$ triplets over n labels such that R is incompatible, but every proper subset of R is compatible. The results given here on the compatibility of triplets appear to have been previously known [20], but are formally proven here.

2 Preliminaries

Given a graph G , we represent the vertices and edges of G by $V(G)$ and $E(G)$ respectively. We use the abbreviated notation uv for an edge $\{u, v\} \in E(G)$. For any $e \in E(G)$, $G - e$ represents the graph obtained from G by deleting edge e . For any integer i , we use $[i]$ to represent the set $\{1, 2, \dots, i\}$.

2.1 Unrooted Phylogenetic Trees

An *unrooted phylogenetic tree* (or just *tree*) is a tree T whose leaves are in one to one correspondence with a label set $L(T)$, and has no vertex of degree two. See Fig. 1(a) for an example. For a collection \mathcal{T} of trees, the *label set* of \mathcal{T} , denoted $L(\mathcal{T})$, is the union of the label sets of the trees in \mathcal{T} . A tree is *binary* if every internal (non-leaf) vertex has degree three. A *quartet* is a binary tree with exactly four leaves. A quartet with label set $\{a, b, c, d\}$ is denoted $ab|cd$ if the path between the leaves labeled a and b does not intersect with the path between the leaves labeled c and d .

For a tree T , and a label set $L \subseteq L(T)$, the *restriction* of T to L , denoted by $T|L$, is the tree obtained from the minimal subtree of T connecting all the leaves with labels in L by suppressing vertices of degree two. See Fig. 1(b) for an example. A tree T *displays* another tree T' , if T' can be obtained from $T|L(T')$ by contracting edges. A tree T displays a collection of trees \mathcal{T} if T displays every tree in \mathcal{T} . If such a tree T exists, then we say that \mathcal{T} is *compatible*; otherwise, we say that \mathcal{T} is *incompatible*. See Fig. 1(a) for an example. Determining if a collection of unrooted trees is compatible is NP-complete [21].

2.2 Multi-State Characters

There is also a notion of compatibility for sets of partitions of a label set L . A *character* χ on L is a partition of L ; the parts of χ are called *states*. If χ has at most r parts, then χ is an r -state character. Given a tree T with $L = L(T)$ and a state s of χ , we denote by $T_s(\chi)$ the minimal subtree of T connecting all leaves




Fig. 1: (a) shows a tree T witnessing that the quartets $q_1 = ab|ce$, $q_2 = cd|bf$, and $q_3 = ad|ef$ are compatible; T is also a witness that the characters $\chi_{q_1} = ab|ce|d|f$, $\chi_{q_2} = cd|bf|a|e$, and $\chi_{q_3} = ad|ef|b|c$ are compatible; (b) shows $T|_{\{a, b, c, d, e\}}$.

with labels having state s for χ . We say that χ is *convex* on T , or equivalently T *displays* χ , if the subtrees $T_i(\chi)$ and $T_j(\chi)$ are vertex disjoint for all states i and j of χ where $i \neq j$. A collection C of characters is *compatible* if there exists a tree T on which every character in C is convex. If no such tree exists, then we say that C is *incompatible*. See Fig. 1(a) for an example. The *perfect phylogeny problem* (or *character compatibility problem*) is to determine whether a given set of characters is compatible.

There is a natural correspondence between quartet compatibility and character compatibility that we now describe. Let Q be a set of quartets, $n = |L(Q)|$, and $r = n - 2$. For each $q = ab|cd \in Q$, we define the r -state character corresponding to q , denoted χ_q , as the character where a and b have state 0 for χ_q ; c and d have state 1 for χ_q ; and, for each $\ell \in L(Q) \setminus \{a, b, c, d\}$, there is a state s of χ_q such that ℓ is the only label with state s for character χ_q (see Example 1). We define the set of r -state characters corresponding to Q by $C_Q = \bigcup_{q \in Q} \{\chi_q\}$.

Example 1. Consider the quartets and characters given in Fig. 1(a): χ_{q_1} is the character corresponding to q_1 , χ_{q_2} is the character corresponding to q_2 , and χ_{q_3} is the character corresponding to q_3 .

The proof of the following lemma relating quartet compatibility to character compatibility is straightforward, and given in the appendix.

Lemma 1. *A set Q of quartets is compatible if and only if C_Q is compatible.*

2.3 Quartet Graphs

We now give a brief overview of quartet graphs which were introduced in [11], and characterize when a collection of quartets is compatible.

Let Q be a collection of quartets with label set L . The *quartet graph* G_Q on Q is the edge-colored graph defined as follows.

1. There is a vertex ℓ in G_Q for each ℓ in L .
2. For every quartet $q = ab|cd$ in Q , G_Q has two edges ab and cd , both labeled by q . We call an edge labeled by q a *q -colored edge*.




Fig. 2: The quartet graph G_Q for the set of quartets given in Fig. 1 is shown in (a). A complete unification sequence for G_Q is shown in (a)–(d). In the figures, the unification of vertices U to a new vertex u is denoted $U \rightarrow u$.

An example of a quartet graph is given in Fig. 2(a). Let G be any edge colored graph. Let $U \subseteq V(G)$ such that for any color c , at most one c -colored edge is incident on the vertices of U . The *unification* of vertices U in G is the graph G' obtained from G as follows:

1. Add a new vertex u to G' .
2. For each c -colored edge vw in G where $v \in U$ and $w \notin U$, add a c -colored edge uw to G' .
3. For each c -colored edge vw in G where both $v \in U$ and $w \in U$, delete all c -colored edges from G' .
4. Delete all vertices in U , and edges incident to vertices in U , from G' .

See Figures 2 and 3 for examples of unification. A *unification sequence* for G is a sequence $G_0=G, G_1, G_2, \dots, G_k$ of graphs where, for any $i > 0$, G_i is derived from G_{i-1} by a unification operation. Note that every graph in a unification sequence that begins with a quartet graph is also a quartet graph. A unification sequence is *complete* if G_k has no edges. See Fig. 2(a)-(d) for an example of a complete unification sequence. The following theorem is from [11].

Theorem 1. *A collection Q of quartets is compatible if and only if there exists a complete unification sequence for the quartet graph G_Q .*

For a quartet graph G , we define the *quartet set corresponding to G* as the set of quartets

$$Q_G = \{ab|cd : \text{there exists edges } ab \text{ and } cd \text{ of the same color in } G\}.$$

3 Compatibility of Quartets

For every $s, t \geq 2$, we fix a set of labels $L_{s,t} = \{a_1, a_2, \dots, a_s, b_1, b_2, \dots, b_t\}$ and define the set

$$Q_{s,t} = \{a_1b_1|a_sb_t\} \cup \bigcup_{i=1}^{s-1} \bigcup_{j=1}^{t-1} \{a_ia_{i+1}|b_jb_{j+1}\}$$

of quartets with label set $L_{s,t}$. We denote the quartet $a_1b_1|a_sb_t$ by q_0 , and a quartet of the form $a_ia_{i+1}|b_jb_{j+1}$ by $q_{i,j}$. See Fig. 3(a) for an illustration.




Fig. 3: The set of quartets $Q_{4,3}$.

Observation 1 For all $s, t \geq 2$, $|Q_{s,t}| = (s-1)(t-1) + 1$.

The proof of the next lemma is straightforward and given in the appendix.

Lemma 2. $Q_{s,t}$ is compatible if and only if $Q_{t,s}$ is compatible.

Lemma 3. For all $s, t \geq 2$, $Q_{s,t}$ is incompatible.

Proof. We use induction on the size of $s+t$.

Base case: Let $s = 2$, $t = 2$ and thus, $s+t = 4$. The set $Q_{2,2}$ contains only the two incompatible quartets $q_0 = a_1b_1|a_2b_2$ and $q_{1,1} = a_1a_2|b_1b_2$. Thus, $Q_{2,2}$ is incompatible.

Induction step: Assume that for every $s' \leq s$ and $t' \leq t$, where $s \geq 2$, $t \geq 2$, $s' \geq 2$, $t' \geq 2$ and $s+t \geq s'+t'+1$, we have that $Q_{s',t'}$ is incompatible. To prove that $Q_{s,t}$ is incompatible, we will show that there is no complete unification sequence for the quartet graph $G = G_{Q_{s,t}}$. For sake of contradiction, assume that there exists a complete unification sequence \mathcal{S} for G . Let U be the set of vertices unified in the first unification operation of \mathcal{S} .

For each $i \in [s]$ and $j \in [t]$, there exists edges of the same color incident on a_i and b_j . Thus, if U contains a vertex a_i , it cannot contain a vertex b_j . Then, w.l.o.g., by Lemma 2, we may assume that U does not contain any vertex b_j for $j \in [t]$. Also, since the quartet $q_0 = a_1b_1|a_sb_t \in Q_{s,t}$, it cannot be the case that U contains both a_1 and a_s . W.l.o.g., we assume that U does not contain a_1 .

Let a_x and a_y be the two vertices in U where x is the smallest index over all of the vertices in U , and y is the largest index over all of the vertices in U . Let G' be the graph resulting from the unification of the vertices of U in G , and let u be the unique vertex in $V(G') \setminus V(G)$. Note that all edges between a_{x-1} and a_x in G become edges between a_{x-1} and u in G' , and all edges between a_y and a_{y+1} in G become edges between u and a_{y+1} in G' . See Fig 3 for an illustration. Let $Q = Q_{G'}$. Since the unification sequence $\mathcal{S} = G, G', G_1, \dots, G_k$ is complete, the sequence G', G_1, \dots, G_k is also complete. Hence, the quartet set Q is compatible.

Let $L = L(Q_{G'})$. Consider the quartet set $Q_{s-(y-x),t}$. Since $|U| > 1$, and, by assumption, $a_1 \notin U$, we have that $1 < x < y \leq s$. Also, by assumption, $t \geq 2$. Hence, $Q_{s-y+x,t}$ is well defined. Let h be an injective mapping from $L_{s-y+x,t}$ to L defined by

$$h(\ell) = \begin{cases} a_i & \text{if } \ell \text{ is of the form } a_i \text{ where } 1 \leq i < x \\ u & \text{if } \ell \text{ is of the form } a_i \text{ where } i = x \\ a_{i+y-x} & \text{if } \ell \text{ is of the form } a_i \text{ where } x < i \leq s-y+x \\ b_i & \text{if } \ell \text{ is of the form } b_i \text{ where } 1 \leq i \leq t \end{cases}$$

We will show that for every quartet $q = \ell_1\ell_2|\ell_3\ell_4 \in Q_{s-y+x,t}$, there exists a quartet $h(\ell_1)h(\ell_2)|h(\ell_3)h(\ell_4) \in Q$. Since $y > x$, $s-y+x+t < s+t$. By the inductive hypothesis, $Q_{s-y+x,t}$ is incompatible. It follows that Q contains an incompatible subset of quartets, contradicting that Q is compatible. We have the following cases.

Case 1: $q = q_0$. Then $\ell_1 = a_1$, $\ell_2 = b_1$, $\ell_3 = a_{s-y+x}$, and $\ell_4 = b_t$. So, $h(\ell_1) = a_1$, $h(\ell_2) = b_1$, and $h(\ell_4) = b_t$. If $y = s$, then $h(\ell_3) = h(a_{s-y+x}) = h(a_x) = u$, and $a_1 b_1 | u b_t \in Q$. If $y < s$, then $h(\ell_3) = h(a_{s-y+x}) = a_{s-y+x+y-x} = a_s$, and $a_1 b_1 | a_s b_t \in Q$.

Case 2: $q = q_{i,j}$ for some $1 \leq i < s - y + x$ and $1 \leq j < t$. Then $\ell_1 = a_i$ and $\ell_2 = a_{i+1}$, $\ell_3 = b_j$, and $\ell_4 = b_{j+1}$. So, $h(\ell_3) = b_j$, and $h(\ell_4) = b_{j+1}$. Note that since $i < s - y + x$, we have that if $i \geq x$, then $y < s$. Hence, we have the following four possibilities to consider.

Case 2a: $1 \leq i < x - 1$. Since both $i < x$ and $i + 1 < x$, $h(\ell_1) = a_i$, $h(\ell_2) = a_{i+1}$, and $a_i a_{i+1} | b_j b_{j+1} \in Q$.

Case 2b: $i = x - 1$. Then $h(\ell_1) = a_{x-1}$, $h(\ell_2) = u$, and $a_{x-1} u | b_j b_{j+1} \in Q$.

Case 2c: $i = x$ and $y < s$. Then, $h(\ell_1) = h(a_x) = u$, $h(\ell_2) = h(a_{x+1}) = a_{x+1+y-x} = a_{y+1}$, and $u a_{y+1} | b_j b_{j+1} \in Q$.

Case 2d: $i > x$ and $y < s$. Then, $h(\ell_1) = h(a_i) = a_{i+y-x}$ and $h(\ell_2) = h(a_{i+1}) = a_{i+1+y-x}$. Since $i > x$, it follows that both $i + y - x > y$ and $i + 1 + y - x > y$. Since $i < s - y + x$, it follows that both $i + y - x < s$ and $i + y - x + 1 < s$. Hence, $a_{i+y-x} a_{i+1+y-x} | b_j b_{j+1} \in Q$.

In every case, we have shown that $h(\ell_1)h(\ell_2)|h(\ell_3)h(\ell_4) \in Q$. \square

Lemma 4. For all $s, t \geq 2$, and every $q \in Q_{s,t}$, $Q_{s,t} \setminus \{q\}$ is compatible.

Proof. Let $q \in Q_{s,t}$. Either $q = q_0$ or $q = q_{x,y}$ for some $1 \leq x < s$ and $1 \leq y < t$. In every case, we present a tree witnessing that $Q_{s,t} \setminus \{q\}$ is compatible.

Case 1. Suppose $q = q_0$. Create the tree T as follows: There is a node for each label in $L_{s,t}$, and two additional nodes a and b . There is an edge ab . For every $a_x \in L_{s,t}$ there is an edge $a_x a$. For every $b_x \in L_{s,t}$, there is an edge $b_x b$. There are no other nodes or edges in T . See Fig. 4(a) for an illustration of T . Consider any quartet $q \in Q_{s,t} \setminus \{q_0\}$. Then $q = a_i a_{i+1} | b_j b_{j+1}$ for some $1 \leq i < s$ and $1 \leq j < t$. Then, the minimal subgraph of T connecting leaves with labels in $\{a_i, a_{i+1}, b_j, b_{j+1}\}$ is the quartet q .

Case 2. Suppose $q = q_{x,y}$ for some $1 \leq x < s$ and $1 \leq y < t$. Create the tree T as follows: There is a node for each label in $L_{s,t}$ and six additional nodes a_ℓ , b_ℓ , ℓ , h , a_h , and b_h . There are edges $a_\ell \ell$, $b_\ell \ell$, ℓh , $h a_h$, and $h b_h$. For every $a_i \in L_{s,t}$, there is an edge $a_i a_\ell$ if $i \leq x$, and an edge $a_i a_h$ if $i > x$. For every $b_j \in L_{s,t}$ there is an edge $b_j b_\ell$ if $j \leq x$, and an edge $b_j b_h$ if $j > y$. There are no other nodes or edges in T . See Fig. 4(b). Now consider any quartet $q \in Q_{s,t} \setminus \{q_{x,y}\}$. Either $q = q_0$ or $q = q_{i,j}$ where $i \neq x$ or $j \neq y$. If $q = q_0$, then the minimal subgraph of T connecting leaves with labels in $\{a_1, b_1, a_s, b_t\}$ is the subtree of T induced by the nodes in $\{a_1, a_\ell, \ell, b_\ell, b_1, a_s, a_h, h, b_h, b_t\}$.




Fig. 4: Illustrating the proof of Lemma 4.

Suppressing all degree two vertices results in a tree that is the same as q_0 . So T displays q . So assume that $q = a_i a_{i+1} | b_j b_{j+1}$ where $i \neq x$ or $j \neq y$. We define the following subset of the nodes in T :

$$V = \begin{cases} \{a_i, a_{i+1}, a_\ell, b_\ell, b_j, b_{j+1}\} & \text{if } i < x \text{ and } j < y, \\ \{a_i, a_{i+1}, a_\ell, b_y, b_\ell, h, b_h, b_{y+1}\} & \text{if } i < x \text{ and } j = y, \\ \{a_i, a_{i+1}, a_\ell, h, b_h, b_j, b_{j+1}\} & \text{if } i < x \text{ and } j > y, \\ \{a_x, a_\ell, h, a_h, a_{x+1}, b_\ell, b_j, b_{j+1}\} & \text{if } i = x \text{ and } j < y, \\ \{a_x, a_\ell, h, a_h, a_{x+1}, b_h, b_j, b_{j+1}\} & \text{if } i = x \text{ and } j > y, \\ \{a_j, a_{j+1}, a_h, h, b_\ell, b_j, b_{j+1}\} & \text{if } i > x \text{ and } j < y, \\ \{a_j, a_{j+1}, a_h, h, b_y, b_\ell, b_h, b_{y+1}\} & \text{if } i > x \text{ and } j = y, \\ \{a_j, a_{j+1}, a_h, h, b_h, b_j, b_{j+1}\} & \text{if } i > x \text{ and } j > y. \end{cases}$$

Now, the subgraph of T induced by the nodes in V is the minimal subgraph of T connecting leaves with labels in q . Suppressing all degree two vertices results in a tree that is the same as q . Hence, T displays q . \square

Theorem 2. For every integer $n \geq 4$, there exists a set Q of quartets over n taxa such that all of the following conditions hold.

1. Q is incompatible.
2. Every proper subset of Q is compatible.
3. $|Q| = \lfloor \frac{n-2}{2} \rfloor \cdot \lceil \frac{n-2}{2} \rceil + 1$.

Proof. By letting $s = \lfloor \frac{n}{2} \rfloor$ and $t = \lceil \frac{n}{2} \rceil$, it follows from Observation 1 and Lemmas 3 and 4 that $Q_{s,t}$ is of size $(\lfloor \frac{n}{2} \rfloor - 1) \cdot (\lceil \frac{n}{2} \rceil - 1) + 1 = \lfloor \frac{n-2}{2} \rfloor \cdot \lceil \frac{n-2}{2} \rceil + 1$ and is incompatible, but every proper subset of $Q_{s,t}$ is compatible. \square

The following theorem allows us to use our result on quartet compatibility to establish a lower bound on $f(r)$.

Theorem 3. Let Q be a set of incompatible quartets over n labels such that every proper subset of Q is compatible, and let $r = n - 2$. Then, there exists a set C of $|Q|$ r -state characters such that C is incompatible, but every proper subset of C is compatible.

Proof. We claim that C_Q is such a set of incompatible r -state characters. Since for two quartets $q_1, q_2 \in Q$, $\chi_{q_1} \neq \chi_{q_2}$, it follows that $|C_Q| = |Q|$. Since Q is incompatible, it follows by Lemma 1 that C_Q is incompatible. Let C' be any proper subset of C . Then, there is a proper subset Q' of Q such that $C' = C_{Q'}$. Then, since Q' is compatible, it follows by Lemma 1 that C' is compatible. \square

Theorem 2 together with Theorem 3 gives the following theorem.

Theorem 4. *For every integer $r \geq 2$, there exists a set C of r -state characters such that all of the following hold.*

1. C is incompatible.
2. Every proper subset of C is compatible.
3. $|C| = \lfloor \frac{r}{2} \rfloor \cdot \lceil \frac{r}{2} \rceil + 1$.

Proof. By Theorem 2 and Observation 1, there exists a set Q of $\lfloor \frac{r}{2} \rfloor \cdot \lceil \frac{r}{2} \rceil + 1$ quartets over $r + 2$ labels that are incompatible, but every proper subset is compatible, namely $Q_{\lfloor \frac{r+2}{2} \rfloor, \lceil \frac{r+2}{2} \rceil}$. The theorem follows from Theorem 3. \square

The quadratic lower bound on $f(r)$ follows from Theorem 4.

Corollary 1. $f(r) \geq \lfloor \frac{r}{2} \rfloor \cdot \lceil \frac{r}{2} \rceil + 1$.

4 Compatibility of Triplets

A *rooted phylogenetic tree* (or just *rooted tree*) is a tree whose leaves are in one to one correspondence with a label set $L(T)$, has a distinguished vertex called the *root*, and no vertex other than root has degree two. See Fig. 5(a) for an example. A rooted tree is *binary* if the root vertex has degree two, and every other internal (non-leaf) vertex has degree three. A *triplet* is a rooted binary tree with exactly three leaves. A triplet with label set $\{a, b, c\}$ is denoted $ab|c$ if the path between the leaves labeled a and b does include the root vertex. For a tree T , and a label set $L \subseteq L(T)$, let T' be the minimal subtree of T connecting all the leaves with labels in L . The *restriction* of T to L , denoted by $T|L$, is the rooted tree obtained from T' by distinguishing the vertex closest to the root of T as the root of T' , and suppressing every vertex other than the root having degree



(a) A tree T witnessing that the triplets $ab|c$, $de|b$, $ef|c$, and $ec|b$ are compatible (b) $T| \{a, b, c, e\}$

Fig. 5: Rooted Phylogenetic Trees

two. A rooted tree T displays another rooted tree T' if T' can be obtained from $T|L(T')$ by contracting edges. A rooted tree T displays a collection of rooted trees \mathcal{T} if T displays every tree in \mathcal{T} . If such a tree T exists, then we say that \mathcal{T} is compatible; otherwise, we say that \mathcal{T} is incompatible. Given a collection of rooted trees \mathcal{T} , it can be determined in polynomial time if \mathcal{T} is compatible [2].

The following theorems follow from the connection between collections of unrooted trees with at least one common label across all the trees, and collections of rooted trees [21].

Theorem 5. *Let Q be a collection of quartets where every quartet in Q shares a common label ℓ . Let R be the set of triplets such that there exists a triplet $ab|c$ in R if and only if there exists a quartet $ab|c\ell$ in Q . Then, Q is compatible if and only if R is compatible.*

Let R be a collection of triplets. For a subset $S \subseteq L(R)$, we define the graph $[R, S]$ as the graph having a vertex for each label in S , and an edge $\{a, b\}$ if and only if $ab|c \in R$ for some $c \in S$. The following theorem is from [4].

Theorem 6. *A collection R of rooted triplets is compatible if and only if $[R, S]$ is not connected for every $S \subseteq L(R)$ with $|S| \geq 3$.*

Corollary 2. *Let R be a set of rooted triplets such that R is incompatible but every proper subset of R is compatible. Then, $[R, L(R)]$ is connected.*

We now contrast our result on quartet compatibility with a result on triplets.

Theorem 7. *For every $n \geq 3$, if R is an incompatible set of triplets over n labels, and $|R| > n - 1$, then some proper subset of R is incompatible.*

Proof. For sake of contradiction, let R be a set of triplets such that R is incompatible, every proper subset of R is compatible, $|L(R)| = n$, and $|R| > n - 1$. The graph $[R, L(R)]$ will contain n vertices and at least n edges. Since each triplet in R is distinct, there will be a cycle C of length at least three in $[R, L(R)]$. Since R is incompatible but every proper subset of R is compatible, by Corollary 2, $[R, L(R)]$ is connected.

Consider any edge e in the cycle C . Let t be the triplet that contributed edge e in $[R, L(R)]$. Let $R' = R \setminus t$. Since the graph $[R, L(R)] - e$ is connected, $[R', L(R')]$ is connected. By theorem 6, R' is incompatible. But $R' \subset R$, contradicting that every proper subset of R is compatible. \square

To show the bound is tight, we first prove a more restricted form of Theorem 2.

Theorem 8. *For every $n \geq 4$, there exists a set of quartets Q with $|L(Q)| = n$, and a label $\ell \in L(Q)$, such that all of the following hold.*

1. Every $q \in Q$ contains a leaf labeled by ℓ .
2. Q is incompatible.
3. Every proper subset of Q is compatible.
4. $|Q| = n - 2$.

Proof. Consider the set of quartets $Q_{2,n-2}$. From Lemmas 3 and 4, $Q_{2,n-2}$ is incompatible but every proper subset of $Q_{2,n-2}$ is compatible. The set $Q_{2,n-2}$ contains exactly $n - 2$ quartets. From the construction, there are two labels in L which are present in all the quartets in $Q_{2,n-2}$. Set one of them to be ℓ . \square

The following is a consequence of Theorem 8 and Theorem 5.

Corollary 3. *For every $n \geq 3$, there exists a set R of triplets with $L(R) = n$ such that all of the following hold.*

1. R is incompatible.
2. Every proper subset of R is compatible.
3. $|R| = n - 1$.

The generalization of the Fitch-Meacham examples given in [16] can also be expressed in terms of triplets. For any $r \geq 2$, let $L = \{a, b_1, b_2, \dots, b_r\}$. Let

$$R_r = ab_r|b_1 \cup \bigcup_{i=1}^{r-1} ab_i|b_{i+1}$$

Let $Q = \{ab|c\ell : ab|c \in R_r\}$ for some label $\ell \notin L$. The set C_Q of r -state characters corresponding to the quartet set Q is exactly the set of characters built for r in [16]. In the partition intersection graph of C_Q , (following the terminology in [16]) labels ℓ and a correspond to the end cliques and the rest of the r labels $\{b_1, b_2, \dots, b_r\}$ correspond to the r tower cliques. From Lemma 1 and Theorem 5, R_r is compatible if and only if Q is compatible.

5 Conclusion

We have shown that for every $r \geq 2$, $f(r) \geq \lfloor \frac{r}{2} \rfloor \cdot \lceil \frac{r}{2} \rceil + 1$, by showing that for every $n \geq 4$, there exists an incompatible set Q of $\lfloor \frac{n-2}{2} \rfloor \cdot \lceil \frac{n-2}{2} \rceil + 1$ quartets over a set of n labels such that every proper subset of Q is compatible. Previous results show that our lower bound on $f(r)$ is tight for $r = 2$ and $r = 3$ [5,7,12,16,17,18]. We give the following conjecture.

Conjecture 2 *For every $r \geq 2$, $f(r) = \lfloor \frac{r}{2} \rfloor \cdot \lceil \frac{r}{2} \rceil + 1$.*

Note that, due to Theorem 3, a proof of Conjecture 2 would also show that the number of incompatible quartets given in the statement of Theorem 2 is also as large as possible. Another direction for future work is to show an upper bound on the function $f(r)$, which would prove Conjecture 1. For quartets, we have a trivial upper bound of $\binom{n}{4}$ on the cardinality of a set of quartets over n labels such that every proper subset is compatible. However, the question for multi-state characters remains open.

Acknowledgments

We thank Sylvain Guillemot and Mike Steel for valuable comments.

References

1. Agarwala, R., Fernández-Baca, D.: A polynomial-time algorithm for the perfect phylogeny problem when the number of character states is fixed. *SIAM J. Comput.* 23(6), 1216–1224 (1994)
2. Aho, A.V., Sagiv, Y., Szymanski, T.G., Ullman, J.D.: Inferring a tree from lowest common ancestors with an application to the optimization of relational expressions. *SIAM J. Comput.* 10(3), 405–421 (1981)
3. Bodlaender, H., Fellows, M., Warnow, T.: Two strikes against perfect phylogeny. In: *Automata, Languages and Programming, Lect. Notes Comput. Sc.*, vol. 623, pp. 273–283. Springer (1992)
4. Bryant, D., Steel, M.: Extension operations on sets of leaf-labelled trees. *Adv. Appl. Math.* 16, 425–453 (1995)
5. Buneman, P.: The recovery of trees from measurements of dissimilarity. In: *Mathematics in the Archeological and Historical Sciences*, pp. 387–395. Edinburgh University Press (1971)
6. Dress, A., Steel, M.: Convex tree realizations of partitions. *Appl. Math. Lett.* 5(3), 3–6 (1992)
7. Estabrook, G.F., Johnson, J., McMorris, F.R.: A mathematical foundation for the analysis of cladistic character compatibility. *Math. Biosci.* 29(1-2), 181–187 (1976)
8. Fernández-Baca, D.: The Perfect Phylogeny Problem. In: *Steiner Trees in Industry*, pp. 203–234. Kluwer (2001)
9. Fitch, W.M.: Toward finding the tree of maximum parsimony. In: *Proceedings of the 8th International Conference on Numerical Taxonomy*. pp. 189–230 (1975)
10. Fitch, W.M.: On the problem of discovering the most parsimonious tree. *The American Naturalist* 111(978), 223–257 (1977)
11. Grünewald, S., Humphries, P.J., Semple, C.: Quartet Compatibility and the Quartet Graph. *Electr. J. Comb.* 15(1), R103 (2008)
12. Gusfield, D.: Efficient algorithms for inferring evolutionary trees. *Networks* 21(1), 19–28 (1991)
13. Habib, M., To, T.H.: On a conjecture of compatibility of multi-states characters. In: *Algorithms in Bioinformatics. Lect. Notes Comput. Sc.*, vol. 6833, pp. 116–127. Springer (2011)
14. Kannan, S., Warnow, T.: Inferring Evolutionary History From DNA Sequences. *SIAM J. Comput.* 23(4), 713–737 (1994)
15. Kannan, S., Warnow, T.: A fast algorithm for the computation and enumeration of perfect phylogenies. *SIAM J. Comput.* 26(6), 1749–1763 (1997)
16. Lam, F., Gusfield, D., Sridhar, S.: Generalizing the Splits Equivalence Theorem and Four Gamete Condition: Perfect Phylogeny on Three-State Characters. *SIAM J. Discrete Math.* 25(3), 1144–1175 (2011)
17. Meacham, C.A.: Theoretical and computational considerations of the compatibility of qualitative taxonomic characters. In: *Numerical Taxonomy, Nato ASI series*, vol. G1. Springer (1983)
18. Semple, C., Steel, M.: *Phylogenetics. Oxford Lecture Ser. Math. Appl.*, Oxford University Press (2003)
19. Shutters, B., Fernández-Baca, D.: A simple characterization of the minimal obstruction sets for three-state perfect phylogenies. *Appl. Math. Lett.* 25(9), 1226–1229 (2012)
20. Steel, M.: Personal communications (2012)
21. Steel, M.: The complexity of reconstructing trees from qualitative characters and subtrees. *J. Classif.* 9(1), 91–116 (1992)

Technical Appendix

5.1 Proof of Lemma 1

Proof. Let T be any tree. It suffices to show that T displays $q = ab|cd$ if and only if χ_q is convex on T . (\Rightarrow) Suppose T displays q . It follows that the path from the leaf labeled a to the leaf labeled b does not intersect the path from the leaf labeled c to the leaf labeled d . Every other state s of χ_q has only one label ℓ_s with state s for χ_q ; hence the minimal subtree connecting all leaves with labels having state s for χ is a single leaf. It follows that χ_q is convex on T . (\Leftarrow) Now suppose that χ_q is convex on T . Then the minimal subtree containing both the leaf labeled a and the leaf labeled b is pairwise disjoint from the minimal subtree containing both the leaf labeled c and the leaf labeled d . It follows that the path from the leaf labeled a to the leaf labeled b does not intersect the path from the leaf labeled c to the leaf labeled d . Hence T displays q . \square

5.2 Proof of Lemma 2

Proof. We define the function $g : L_{s,t} \rightarrow L_{t,s}$ by

$$g(\ell) = \begin{cases} b_i & \text{if } \ell \text{ is of the form } a_i \text{ for some } 1 \leq i \leq s \\ a_i & \text{if } \ell \text{ is of the form } b_i \text{ for some } 1 \leq i \leq t. \end{cases}$$

It is straightforward to verify that $\ell_1\ell_2|\ell_3\ell_4 \in Q_{s,t}$ if and only if $g(\ell_1)g(\ell_2)|g(\ell_3)g(\ell_4) \in Q_{t,s}$. The theorem follows. \square